

PATIENT INSTRUCTIONS GLUCOSE TOLERANCE TESTS

- 1. Fast for 10- 12 hours before the test (NEVER FAST OVER 16 HOURS!)
- 2. You may have water
- 3. You may have essential medications
- 4. If you breast feed express milk before the test begins or feed your baby before the test begins. You cannot breast feed during the test.
- 5. You can make an appointment for the test although this is not essential.
- 6. A glucose tolerance test for suspected hyperglycaemia will take 2 hours.
- 7. A glucose tolerance test + insulin's will take 2 hours & you MUST fast for at least 12 hours.
- 8. A glucose tolerance test for hypoglycaemia will take 5 hours.

WHAT TO EXPECT

- 1. You will have your height & weight measured & this will require that you remove your shoes.
- 2. A fasting sample of blood will be collected from you.
- 3. You will then be given a 300ml bottle of glucose to drink (similar to lemonade).
- 4. You will be asked to drink the complete bottle of drink within 5 minutes.
- A clock with an alarm will be started as soon as you drink the last mouthful of glucose & this
 will be set for 2 hours (for suspected hyperglycaemia) or 1 hour (for suspected
 hypoglycaemia)
- 6. YOU WILL BE REQUIRED TO REST DURING THIS WAITING PERIOD & NOT EAT OR SMOKE.
- 7. You may drink water only.
- 8. When the alarm sounds you will have another sample of blood taken.
- 9. If you are being tested for suspected hypoglycaemia you will have hourly blood samples taken for the 5 hours duration of the test.
- 10. If you are having insulin levels taken with your glucose tolerance test you will have your blood taken hourly.
- 11. Remember to bring you reading glasses if you use them because it is essential that you check your name on each sample collected.
- 12. You can expect your results to be back at your doctor's surgery within 2 working days.